



YEAR 2010

[Metabolic endotoxemia and saturated fat contribute to circulating NGAL concentrations in subjects with insulin resistance.](#) Moreno-Navarrete JM, Manco M et al. Int J Obes (Lond). 2010 Feb;34(2):240-9. doi: 10.1038/ijo.2009.242.

[Relation between human LPIN1, hypoxia and endoplasmic reticulum stress genes in subcutaneous and visceral adipose tissue.](#) Miranda M, Escoté X et al. Int J Obes (Lond). 2010 Apr;34(4):679-86. doi: 10.1038/ijo.2009.290.

[Circulating bactericidal/permeability-increasing protein \(BPI\) is associated with serum lipids and endothelial function.](#) Esteve E, Castro A et al. Thromb Haemost. 2010 Feb 19;103(4):780-7. doi: 10.1160/TH09-08-0596.

[Tumor Necrosis-Like Weak Inducer of Apoptosis as a Proinflammatory Cytokine in Human Adipocyte Cells: Up-Regulation in Severe Obesity Is Mediated by Inflammation But Not Hypoxia.](#) Vendrell J, Maymó-Masip E et al. J Clin Endocrinol Metab. 2010 Jun;95(6):2983-92. doi: 10.1210/jc.2009-2481.

[Retinol-binding protein 4 levels in obese children and adolescents with glucose intolerance.](#) Yeste D, Vendrell J et al. Horm Res Paediatr. 2010;73(5):335-40. doi: 10.1159/000308165.

[PP2A regulatory subunit PP2A-B' counteracts S6K phosphorylation.](#) Hahn K, Miranda M et al. Cell Metab. 2010 May 5;11(5):438-44. doi: 10.1016/j.cmet.2010.03.015.

[Paired Subcutaneous and Visceral Adipose Tissue Aquaporin-7 Expression in Human Obesity and Type 2 Diabetes: Differences and Similarities between Depots.](#) Miranda M, Escoté X et al. J Clin Endocrinol Metab. 2010 Jul;95(7):3470-9. doi: 10.1210/jc.2009-2655.

[Obesity and Insulin Resistance-Related Changes in the Expression of Lipogenic and Lipolytic Genes in Morbidly Obese Subjects.](#) Tinahones FJ, Garrido-Sánchez L et al. Obes Surg. 2010 Nov;20(11):1559-67. doi: 10.1007/s11695-010-0194-z.

[Lpin1 in human visceral and subcutaneous adipose tissue: similar levels but different associations with lipogenic and lipolytic genes.](#) Miranda M, Escoté X et al. Am J Physiol Endocrinol Metab. 2010 Aug;299(2):E308-17. doi: 10.1152/ajpendo.00699.2009.

[Cyclin G2 regulates adipogenesis through PPAR gamma coactivation.](#) Aguilar V, Annicotte JS et al. Endocrinology. 2010 Nov;151(11):5247-54. doi: 10.1210/en.2010-0461.

[Weight loss in prepubertal obese children is associated with a decrease in adipocyte fatty acid-binding protein without changes in lipocalcin - 2: a two-year longitudinal study.](#) Corripio R, González-Clemente JM A et al. Eur J Endocrinol. 2010 Dec;163(6):887-93. doi: 10.1530/EJE-10-0408.



DIABETES AND METABOLIC
ASSOCIATED DISEASES
RESEARCH GROUP

Is plasma 25(OH) D related to adipokines, inflammatory cytokines and insulin resistance in both a healthy and morbidly obese populations? Vilarrasa N, Vendrell J et al. Endocrine. 2010 Oct;38(2):235-42. doi: 10.1007/s12020-010-9379-4.